CONTENTS

Vol. 154, No. 1, January 1988

Effects of human pancreatic polypeptide on exocrine pancreatic function in conscious rats. *A. Funakoshi, K. Miyasaka, H. Shinozaki, K. Kitani, N. Sugiyama and H. Yajima* 1

The relationship between cholelithiasis and diabetes mellitus: Discussion of age, obesity, hyperlipidemia and neuropathy. *N. Tanno, M. Koizumi and Y. Goto* 11

Extraadrenal immunohistochemical distribution of steroid 21-hydroxylase in human. *H. Sasano and N. Sasano* 21

Intravenous insulin infusion system to determine insulin regimen in diabetics. *M. Ito, T. Sato, T. Saito, M. Kokubun, M. Inoue, K. Saito and K. Yoshinaga* 29

Evaluation of monoclonal antibody against human pulmonary surfactant apoprotein. *K. Takeda, I. Mizusawa, Y. Kuroda and K. Hiraiwa* 37

Changes in the motility pattern of human spermatozoa during in vitro incubation. *K. Hoshi, K. Yanagida, T. Aita, N. Yoshimatsu and A. Sato* 47

Rods also participate in human color vision. *Y. Sugita and K. Tasaki* 57

Angiogenesis factors in ocular tissues of normal rabbits on chorioallantoic membrane assay. *T. Okamoto, S. Oikawa, T. Toyota and Y. Goto* 63

The irradiation effects on the cytoskeletons of C3H/He mouse mammary tumor cells and vascular basement membrane in relation to vascular invasion: A model of intraoperative radiotherapy. *M. Kobayashi* 71

Short Reports

A new treatment for obstructive sleep apnea syndrome by electrical stimulation of submental region. *H. Miki, W. Hida, H. Inoue and T. Takishima* 91

Complete prevention of acetylcholine induced coronary artery spasm with nifedipine. *K. Shimizu, K. B. Ogawa, T. Kato, J.-D. Lee, A. Hara and T. Nakamura* 93

Vol. 154, No. 2, February 1988

Fibronectin localization in the mouse embryo from the two cell stage to the morula stage. *T. Yohkaichiya, H. Hoshiai, S. Uehara and A. Yajima* 95

Responsiveness of peripheral blood lymphocytes from cancer patients and healthy donors to interleukin-2 (IL-2). *M. Kambe, Y. Mitachi, R. Kanamaru and A. Wakui* 101

The relationship between walking speed and muscle strength for knee extension in hemiparetic stroke patients: A follow-up study. *R. Nakamura, S. Watanabe, T. Handa and I. Morohashi* 111

Properties of thyrotropin (TSH) binding to plasma membranes prepared from rat fat tissues. *K. Kaise, N. Kaise, K. Yoshida, Y. Itagaki, Y. Kiso, T. Arai, M. Yamamoto, T. Sakurada, S. Saito and K. Yoshinaga* 115

Pressor response to vasopressin and norepinephrine in DOC-salt hypertensive and prehypertensive rats. *Y. Ouchi, Y. Yazaki, R.-C. Tsai and T. Ashida* 125

Endoscopic diagnosis and treatment of Dieulafoy's ulcer. *S. Asaki, H. Sato, T. Nishimura, S. Ohkubo, R. Yamagata, S. Ito, Y. Saito, S. Miyazaki and N.*

Yaginuma 135

Chronic granulomatous disease with neutrophil membrane cytochrome b deficiency: Demonstration by immunochemical staining with monoclonal antibody. *N. Minegishi, M. Nakamura, K. Suzaki, M. Terasawa, M. Minegishi, S. Tsuchiya and T. Konno* 143

Persistent jaundice after hepatic portojejunostomy in biliary atresia: Are the patients' prognoses determined within 3 months after surgery? *T. Chiba, R. Ohi, T. Uchida, Y. Hayashi and I. Mochizuki* 149

Comparative analysis of lymphocyte phenotypes between carriers of human immunodeficiency virus (HIV) and adult patients with primary immunodeficiency using two-color immunofluorescence flow cytometry. *T. Ohno, T. Kanoh, T. Suzuki, T. Masuda, K. Kuribayashi, S. Araya, H. Arai and H. Uchino* 157

The hypotensive mechanism of percutaneous transluminal dilatation (PTD) in renovascular hypertension due to bilateral renal artery stenosis. *S. Sasaki, Y. Imai, K. Abe, M. Nihei, N. Minami, M. Munakata, H. Sasaki, H. Sekino and K. Yoshinaga* 173

Role of calcium in suppression of glucagon release from isolated diabetic rat pancreata. *Y. Fujita, K. Kawaji, T. Moriya, K. Matoba, K. Ishii, Y. Yajima and H. Okabe* 185

Induction of the $CD4^-8^+$ suppressor phenotype in $CD4^+8^+$ human thymocytes by phorbol myristate acetate. *Y. Sakiyama, A. Ishizaka, T. Watanabe, T. Ariga and S. Matsumoto* 195

Plasma levels of atrial natriuretic peptide in patients with borderline and essential hypertension. *T. Nakamura, S. Ichikawa, T. Sakamaki, M. Fujie, A. Yagi, T. Kurashina and K. Murata* 205

Vol. 154, No. 3, March 1988

Roentgenographically occult bronchogenic carcinoma of non-polypoid type: Histological patterns of longitudinal growth within the bronchus. *N. Nagamoto, Y. Saito, H. Suda, T. Imai, M. Sato, S. Ohta, K. Kanma, M. Sagawa, S. Takahashi, K. Usuda, T. Nakada, H. Sato and K. Hashimoto* 215

Effect of tetrahydrobiopterin and 5-hydroxytryptophan on hereditary progressive dystonia with marked diurnal fluctuation: A suggestion of the serotonergic system involvement. *A. Ishida, G. Takada, Y. Kobayashi, I. Toyoshima and K. Takai* 233

Walking cycle after stroke. *R. Nakamura, T. Handa, S. Watanabe and I. Morohashi* 241

Human cytotrophoblastic antigens defined by monoclonal antibodies. *Y. Abe, K. Okamura, Y. Hamazaki, Y. Wada and A. Yajima* 245

Effects of androgen on 17β-estradiol production by cultured human granulosa cells. *Y. Wada, A. Tsuiki, T. Fukaya, O. Shinkawa, S. Satoh, M. Horiguchi, H. Hoshiai and A. Yajima* 253

Fibronectin localization in the mouse embryo from the blastocyst stage to the egg cylinder stage in vitro. *T. Yohkaichiya, H. Hoshiai and A. Yajima* 261

Blood vascular organization of the human appendix: A scanning electron microscopic study of corrosion casts. *S. Hashimoto and T. Ogata* 271

Ultrastructural abnormalities in the skeletal muscle of children with chronic

cholestasis and a long-term vitamin E replacement. *Y. Kobayashi, Y. Tazawa, M. Nakagawa, O. Higashi and T.Y. Yamamoto* 285
Apolipoprotein E phenotypes of normo- and hyperlipoproteinemia in Japanese. *R. Sano, R. Abe, S. Oikawa, Y. Fujii, S. Hori, N. Suzuki, T. Toyota and Y. Goto* 297
Kinetics of mixed venous oxygen tension during rhythmic contraction of the hindlimb of dogs. *T. Kutsuzawa and H. Yamabayashi* 305

Vol. 154, No. 4, April 1988

Increased plasma concentrations of complement modulating proteins (C1 inhibitor, C4-binding protein, factor H and factor I) in psoriasis. *K. Ohkohchi, W. Torinuki and H. Tagami* 315
Accuracy of digital palpation for size assessment of the prostate evaluated by transrectal sonography. *H. Ohe, K. Ohnishi, H. Watanabe, Y. Itakura and T. Inaba* 323
Effects of calcium antagonists on low density lipoprotein metabolism in human arterial smooth muscle cells. *S. Mori, H. Ito and K. Yamamoto* 329
Natural killer cells in dilated cardiomyopathy. *A. Yokoyama* 335
Highly efficient procedure for production of human monoclonal antibodies: Establishment of hybrids between Epstein-Barr virus-transformed B lymphocytes and heteromyeloma cells by use of GIT culture medium. *T. Kudo, A. Asao and T. Tachibana* 345
Physiological significance of fetal blood gas changes elicited by different delivery postures. *S. Koga, Y. Koga and H. Nagai* 357
A new bleeding tendency due to hereditary hyper α_2-macroglobulinemia. *Y. Endo, K. Iwamoto, S. Mamiya, H. Niitsu, T. Itoh and A.B. Miura* 365
Effect of warning signal on reaction time and EMG activity of the biceps brachii muscle in elbow flexion and forearm supination. *J.A.P. Mojica, Y. Yamada and R. Nakamura* 375
Twenty four-hour urinary excretion of sodium, potassium and urea in Japanese children. *M. Moriyama and H. Saito* 381
A case-control study of lung cancer in nonsmoking women. *H. Shimizu, M. Morishita, K. Mizuno, T. Masuda, Y. Ogura, M. Santo, M. Nishimura, K. Kunishima, K. Karasawa, K. Nishiwaki, M. Yamamoto, S. Hisamichi and S. Tominaga* 389
Cyclic GMP accumulation by atriopeptins in cultured rat renal and vascular smooth muscle cells. *M. Sato, K. Abe, K. Takeuchi, K. Matsui, M. Yasujima, S. Fang, M. Kanazawa, K. Yoshida, T. Kimura and K. Yoshinaga* 399

Short Reports

Does diffuse intimal thickening in human coronary artery act as a diffusion barrier to endothelium-derived relaxing factor? *M. Keitoku, H. Okayama, Y. Satoh, Y. Maruyama and T. Takishima* 413
The receptor mechanisms for histamine actions in proximal portion differ from those in distal portion in human coronary artery. *M. Keitoku, Y. Maruyama and T. Takishima* 415